

Euvax B inj.

HEPATITIS B VACCINE, RECOMBINANT

WHO U'la

Euvax B consists of highly purified, non-infectious particles of Hepatitis B surface antigen (HBsAg) adsorbed onto aluminum salts as an adjuvant and preserved with thimerosal. It is a recombinant DNA hepatitis B vaccine derived from HBsAg produced by DNA recombinant technology in yeast cells (*Saccharomyces cerevisiae*). The vaccine meets the WHO requirements for recombinant hepatitis B vaccines. No substances of human origin are used in its manufacture.

DESCRIPTION

Euvax B is a white, slightly opalescent suspension.

COMPOSITION

1 ml of the above vaccine contains:

- Active ingredient: Purified HBsAg----- 20 µg
 - Adjuvant: Aluminum Hydroxide Gel (as aluminum) ----- 0.5 mg
 - Preservative: Thimerosal----- 0.01 w/v%
- Excipients: Potassium phosphate, monobasic, Sodium phosphate, dibasic, Sodium chloride

INDICATION AND USAGE

Immunization against infection caused by all known subtypes of Hepatitis B virus.

DOSAGE AND ADMINISTRATION

Euvax B should be injected intramuscularly into the deltoid muscle of older children or adults.

- One adult dose (from 16 years) is 1.0 ml containing 20 µg of HBsAg.

The immunization regimen consists of three doses of vaccine given according to the following schedule:

- 1 st dose: at elected date
- 2nd dose: 1 month after the first dose
- 3rd dose: 6 month after the first dose

Booster vaccination: the WHO does not recommend booster vaccination, as it has been shown that 3 dose series of hepatitis B immunization protects for as long as 15 years, and that a protective anamnestic response occurs after exposure to HBV, even if protective antibodies have been lost over time. However, some local vaccination programmes worldwide currently include a recommendation for a booster dose, and these should be respected.

An alternative 0-, 1-, and 2-month schedule and a 12-month booster may be used in certain populations (e.g., neonates born from Hepatitis B infected mothers, someone who has or might have been recently exposed to the virus, or certain travelers to high-risk areas). Additional dose(s) of vaccine may be required in hemodialysis or immunodeficient patients, since protective antibody titers ($> 10 \text{ IU/l}$) may not be obtained after the primary immunization course.

CONTRAINDICATIONS

Hepatitis B vaccine is contraindicated for use in persons with hypersensitivity to any component of Euvax B. The vaccine will not harm individuals currently or previously infected with HB virus. Individuals infected with human immunodeficiency virus (HIV), both asymptomatic and symptomatic, should be immunized with hepatitis B vaccine according to standard schedules.

WARNINGS AND PRECAUTIONS

General precautions:

- The administration of Euvax B should be postponed in patients suffering from acute, severe febrile illness.

- In patients suffering from multiple sclerosis, any stimulation of the immune system can induce exacerbation of their symptoms. Therefore, for these patients the benefits of vaccination against Hepatitis B should be weighed against the risks of exacerbation of multiple sclerosis.

- As with all injectable reactions following the administration of the vaccine. B should be weighed that protection cannot be obtained by vaccination in patients in latent or progressive state of Hepatitis B.

- It is considered that protection should always be readily available in case of rare anaphylactic reactions following the administration of the vaccine.

- As with all injectable reactions following the administration of the vaccine.

Precautions for use:

- Shake before administration, since a fine white deposit with a clear colorless supernatant may form during storage.

- A sterile syringe and sterile needle should be used for each injection. However, as with all inactivated

- Pregnancy and lactation Euvax B should be used during fetal development has not been assessed. Euvax B should be used during

- The effect of the HBsAg on fetal development has not been assessed. Euvax B should be used during

- The effect of the HBsAg on fetal development has not been assessed. Euvax B should be used during

- The effect of the HBsAg on fetal development has not been assessed. Euvax B should be used during

- The effect of the HBsAg on fetal development has not been assessed. Euvax B should be used during

- The effect of the HBsAg on fetal development has not been assessed. Euvax B should be used during

- The effect of the HBsAg on fetal development has not been assessed. Euvax B should be used during

ADVERSE REACTIONS

Gastrointestinal disorders

Rare: nausea

Common: abdominal pain, diarrhea, vomiting

General disorders and administration site conditions

Rare: malaise, fatigue

Common: fever, induration, oedema, tenderness, inflammation

Very common: injection site pain

Infections and infestations

Uncommon: moniliasis, rhinitis

Investigations

Rare: transient increase of transaminase

Metabolism and nutrition disorders

Common: anorexia

Musculoskeletal and connective tissue disorders

Rare: myalgia, arthritis

Nervous system disorders

Very rare: optic neuritis, facial paralysis, Guillain-Barre syndrome, aggravation of disseminated

sclerosis

Rare: headache, dizziness

Common: crying abnormal, and perinatal conditions

Pregnancy, puerperium and perinatal conditions

Uncommon: jaundice neonatal

Psychiatric disorders

Common: insomnia, nervousness, irritability

Skin and subcutaneous tissue disorders

Common: rash erythematous, erythema

Common: rash erythematous, rash, rash maculo-papular


Uncommon: pityriasis rosea, rash, rash maculo-papular

Vascular disorders

Common: hematoma

STORAGE CONDITIONS

Do not exceed the expiry date stated on the external packaging.
Store between 2°C and 8°C (in a refrigerator). Do not freeze.



PRESENTATION 1ml/vial x 20 vials - 1 ml/vial x 10 vials - 1 ml/vial x 1 vial

Issuance date: Jan. 26, 2017

Revised date: Nov. 15, 2019

Manufactured by: LG Chem, [Factory 1] 129, Seokam-ro, Iksan-si, Jeollabuk-do, Korea,
[Factory 2] 151, Osongsaengmyeong 1-ro, Osong-eup, Heungdeok-gu, Cheongju-si, Chungcheongbuk-
do, Korea.